


Current Biology

The Astrin-SKAP complex reduces friction at the kinetochore-microtubule interface

Graphical abstract



Authors

Miquel Rosas-Salvans,
Renaldo Sutanto, Pooja Suresh,
Sophie Dumont

Correspondence

miquel.rosas377@gmail.com (M.R.-S.),
sophie.dumont@ucsf.edu (S.D.)

In brief

Rosas-Salvans et al. show that the Astrin-SKAP complex reduces friction at the mammalian kinetochore-microtubule interface, in contrast to other kinetochore proteins thought to increase friction. Astrin-SKAP increases interface dynamics and force responsiveness, effectively lubricating attachments as they stabilize, which may help preserve them.

Highlights

- SKAP increases kinetochore motility during mammalian metaphase
- SKAP decreases friction at the kinetochore-microtubule interface, lubricating it
- SKAP increases kinetochore-microtubule interface's dynamics and force responsiveness
- Reducing interface friction independently of grip may be key to robust attachments

Article

The Astrin-SKAP complex reduces friction at the kinetochore-microtubule interface

Miquel Rosas-Salvans,^{1,5,*} Renaldo Sutanto,¹ Pooja Suresh,^{1,2} and Sophie Dumont^{1,2,3,4,6,7,*}

¹Department of Bioengineering & Therapeutic Sciences, UCSF, 600 16th Street, San Francisco, CA 94158, USA

²Biophysics Graduate Program, UCSF, 600 16th Street, San Francisco, CA 94158, USA

³Department of Biochemistry & Biophysics, UCSF, 600 16th Street, San Francisco, CA 94158, USA

⁴Chan Zuckerberg Biohub, 499 Illinois Street, San Francisco, CA 94158, USA

⁵Twitter: @RosasSalvans

⁶Twitter: @DumontLab

⁷Lead contact

*Correspondence: miquel.rosas377@gmail.com (M.R.-S.), sophie.dumont@ucsf.edu (S.D.)

<https://doi.org/10.1016/j.cub.2022.04.061>

SUMMARY

The kinetochore links chromosomes to spindle microtubules to drive chromosome segregation at cell division. While we know nearly all mammalian kinetochore proteins, how these give rise to the strong yet dynamic microtubule attachments required for function remains poorly understood. Here, we focus on the Astrin-SKAP complex, which localizes to bioriented kinetochores and is essential for chromosome segregation but whose mechanical role is unclear. Live imaging reveals that SKAP depletion dampens the movement and decreases the coordination of metaphase sister kinetochores and increases the tension between them. Using laser ablation to isolate kinetochores bound to polymerizing versus depolymerizing microtubules, we show that without SKAP, kinetochores move slower on both polymerizing and depolymerizing microtubules and that more force is needed to rescue microtubules to polymerize. Thus, in contrast to the previously described kinetochore proteins that increase the grip on microtubules under force, Astrin-SKAP reduces the grip, increasing attachment dynamics and force responsiveness and reducing friction. Together, our findings suggest a model where the Astrin-SKAP complex effectively “lubricates” correct, bioriented attachments to help preserve them.

INTRODUCTION

The kinetochore links each chromosome to spindle microtubules at cell division, transmitting spindle forces to move chromosomes. To perform its function, the kinetochore must not only bind microtubules strongly enough to resist cellular forces but also slide on them to move and segregate chromosomes. Although we now have a detailed map of mammalian kinetochore components and are uncovering their structure, biochemistry, and biophysics, how these components together give rise to the mechanics of the kinetochore-microtubule interface remains poorly understood. Indeed, we cannot as yet reconstitute mammalian kinetochores or the microtubule bundles they bind to *in vitro*, and applying precise mechanical perturbations to mammalian kinetochores remains challenging *in vivo*. How mammalian kinetochore-microtubule attachments can be robust and strong yet dynamic remains an open question. Answering this question is central to understanding how cells accurately segregate their chromosomes.

To perform its function, the kinetochore-microtubule interface both generates and responds to force. In mammalian cells, the kinetochores bind to the 15–25 microtubules that form a kinetochore-fiber (k-fiber)^{1,2} and that both polymerize and

depolymerize. When sister kinetochores oscillate together at metaphase, active (energy-consuming) force generation from the microtubule depolymerization at the “front” kinetochore (moving toward the pole) largely drives the movement of the pair; in turn, passive, frictional force at the “back” kinetochore (moving away from the pole) is generated as the kinetochore proteins slide on microtubules and oppose movement (Figure 1A).^{3–8}

The kinetochore-microtubule interface also responds to force. For example, force coordinates microtubule dynamics at both sister kinetochores as chromosomes move^{7,9,10} and helps maintain chromosomes in the spindle center through spatially regulated polar ejection forces.^{11,12} Key to the interface’s ability to generate and respond to force is that it is dynamic: this allows kinetochore mobility on microtubules and microtubule growth and shrinkage, and as such perturbing microtubule dynamics causes segregation defects.¹³ While we now know different kinetochore molecules that generate force and increase grip at the microtubule interface,^{14–21} the mechanisms that make this interface dynamic and able to respond to force despite this grip remain poorly defined.

Ndc80 and Ska kinetochore complexes play central roles in microtubule attachment in mammalian cells. The Ndc80 complex is essential to the formation of kinetochore-microtubule attachments *in vivo*¹⁴ and directly binds microtubules, forming




Figure 1. SKAP increases kinetochore mobility and is essential for sister kinetochore coordination in metaphase

(A) Simplified representation of metaphase chromosome oscillations. The force from depolymerizing microtubules (purple) at the front kinetochore drives the movement of both sisters. Frictional force at the back kinetochore, which is bound to polymerizing microtubules (pink), opposes movement.

(B) Representative live images (left) of control and siSKAP metaphase Rpe1-GFP cells (GFP-CenpA and centrin1-GFP), with red boxes highlighting the regions used for kymographs (right) of centriole and kinetochore movements for those cells. Scale bars, 10 μm or 1 μm .

(C) The standard deviation of the position of individual control and siSKAP metaphase kinetochores over time (Mann-Whitney test).

(D) The average speed of individual control and siSKAP metaphase kinetochores (Mann-Whitney test).

(E) Velocity correlation between metaphase sister kinetochores (Mann-Whitney test).

(F) Fraction of time in which individual metaphase sister kinetochores move in opposite directions (Mann-Whitney test).

(legend continued on next page)

load-bearing attachments *in vitro*.^{22,23} During mitosis, progressive dephosphorylation of the Ndc80 component Hec1 increases its microtubule affinity,²⁴ load-bearing ability,²⁰ and grip on polymerizing microtubules.^{17,18} Thus, Ndc80 binds more stably to microtubules as attachments mature, with one phosphorylation site (S69 on protein Hec1) maintaining basal interface dynamics.²⁵ In turn, the Ska complex is essential for proper chromosome alignment and mitotic progression but is only loaded at kinetochores once they biorient and in an Ndc80-dependent manner.²⁶ *In vivo*, Ska increases attachment stability to depolymerizing microtubules under force.¹⁹ *In vitro*, Ska directly binds microtubules²⁷ and increases Ndc80's affinity for microtubules and microtubule tracking and load-bearing ability on depolymerizing microtubules.^{16,20,28} Thus, Ska is thought to be a "locking" factor increasing the grip on microtubules and stabilizing mature attachments. In addition to Ndc80 and Ska, the Astrin-SKAP (SKAP for short hereafter) complex has been proposed to contribute to microtubule attachment. Like Ska, SKAP is essential for chromosome alignment and mitotic progression and is only loaded at bioriented kinetochores in a Ndc80-dependent manner.^{29–31} Strikingly, similar to Ndc80 dephosphorylation,^{18,32} SKAP depletion decreases k-fiber poleward flux,³³ suggesting that SKAP's presence at the kinetochore may not increase the grip on microtubules. Yet SKAP directly interacts with microtubules *in vitro*,^{34,35} synergistically with Ndc80.³⁵ Similar to Ndc80 and Ska,^{36,37} mutations or changes in SKAP expression are highly frequent in some cancers and increase aneuploidy.^{38,39} Yet SKAP's mechanical role at the kinetochore-microtubule interface, if any, is not known. More broadly, whether all microtubule-binding kinetochore proteins increase microtubule grip, as Ndc80 and Ska complexes do, or whether the presence of some proteins instead reduce the grip to "lubricate" the attachment, remains an open question.

Here, we show that SKAP decreases the grip at the kinetochore-microtubule interface, effectively lubricating it. We use live imaging to show that SKAP increases the magnitude of metaphase kinetochore movements and the coordination between sisters and that it decreases the tension the sisters are under. Using laser ablation, we show that SKAP increases the velocity with which kinetochores move on both polymerizing and depolymerizing microtubules and that it makes the dynamics of attached microtubules more force responsive. Thus, not all kinetochore proteins increase the grip on microtubules: SKAP does exactly the opposite, reducing the friction at the interface. We propose that SKAP promotes accurate segregation by lubricating correct, mature attachments, ensuring that they can smoothly slide despite the mechanisms that stabilize them late in mitosis. More broadly, our work suggests that maintaining a strong yet dynamic kinetochore-microtubule interface not only requires components that grip—which are being actively studied—but also components that help slide, which are distinct from those that grip.

RESULTS

SKAP increases kinetochore mobility and is essential for sister kinetochore coordination in metaphase

To probe the mechanical role of the Astrin-SKAP complex at the kinetochore-microtubule interface, we live imaged metaphase chromosome movements in human retinal pigment epithelial-1 (human Rpe1), GFP-CenpA (kinetochore), and centrin1-GFP (centriole) cells (Rpe1-GFP⁴⁰ cells hereafter). Upon SKAP depletion by RNAi (siSKAP; **Figures S1A** and **S1B**), kinetochore pairs oscillated less far from their mean position than the control ($0.4 \pm 0.2 \mu\text{m}$ versus $0.6 \pm 0.2 \mu\text{m}$ SD; **Figures 1B** and **1C**; **Videos S1** and **S2**), as expected.³³ siSKAP kinetochores moved at a slower velocity than the control (1.2 ± 0.4 versus $1.8 \pm 0.4 \mu\text{m}/\text{min}$; **Figures 1B** and **1D**; **Videos S1** and **S2**), indicating that SKAP increases kinetochore mobility. This suggests that SKAP increases the kinetochore-microtubule interface dynamics. Additionally, the sister kinetochore movement coordination decreased without SKAP. siSKAP sister kinetochore pairs showed a lower velocity correlation than the control sister pairs (0.54 ± 0.18 versus 0.74 ± 0.12 ; **Figure 1E**), with siSKAP sister kinetochores moving in opposite directions for a higher fraction of time than the control ($29\% \pm 7\%$ versus $17\% \pm 7\%$; **Figure 1F**). SKAP silencing using an alternative siRNA sequence confirmed the specificity of the oscillation phenotypes associated with SKAP silencing (**Figures S1C–S1G**). Furthermore, while in the control sister kinetochores, the front kinetochore usually reverses direction before the back kinetochore (**Figures 1G** and **S1H**),⁷ in siSKAP sisters, this preference was lost and the back kinetochore switched first more often than the control (41% versus 19% ; **Figure 1G**). Together, these findings indicate that SKAP is essential for sister kinetochore mobility and coordination at metaphase.

SKAP decreases tension at the kinetochore-microtubule interface

The mechanical force from the sister kinetochore and from the spindle on chromosome arms is thought to coordinate sister kinetochore movement and inform directional kinetochore switching.^{3,7,12} To probe whether the miscoordination in the siSKAP sister kinetochore movement (**Figures 1E–1G**) could be due to defects in force generation or in how the kinetochore respond to force, we measured the interkinetochore (K-K) distance in siSKAP cells. Decreasing the activity of the microtubule-binding kinetochore proteins (such as Hec1 phosphorylation or Ska depletion) typically reduces the kinetochore's grip or ability to sustain force before sliding on microtubules and thus leads to a lower K-K distance^{17,41,42} (**Figure 2A**). By contrast, we found that in live Rpe1-GFP cells, siSKAP kinetochore pairs had a higher K-K distance than the control ($1.4 \pm 0.2 \mu\text{m}$ versus $1.1 \pm 0.1 \mu\text{m}$; **Figure 2B**), consistent with some³⁰ but not other^{29,43} previous reports. We used an alternative siRNA sequence to silence SKAP and confirmed that the increase in

(G) Fraction of metaphase directional switches in which the front or back kinetochore switches first or both switch together (Fisher's exact test) (n_s = number of switches).

(C)–(G) are from individual kinetochore tracks obtained from the dataset in (B) (n = number of kinetochore pairs, 1–4 kinetochore pairs per analyzed cell from 18 control and 20 siSKAP cells).

See also **Figure S1** and **Videos S1** and **S2**.




Figure 2. SKAP decreases tension at the kinetochore-microtubule interface

(A) High tension between sister kinetochores leads to a high K-K distance (red double arrow). High tension can stem from high spindles forces, a tighter grip of kinetochores on spindle microtubules, or both.

(B) K-K distance average over time for individual kinetochore pairs in the control and siSKAP Rpe1-GFP cells from the dataset in Figure 1 (Student's t test) (n = number of kinetochore pairs, 1–4 kinetochore pairs per analyzed cell from 18 control and 20 siSKAP cells).

(C) The time that the individual control or siSKAP cells spend from nuclear envelope breakdown (NEB) to anaphase onset (Mann-Whitney test) (n = number of cells). Box and whiskers graph (quartiles including 5%–95%).

(D) Representative immunofluorescence images in the control and siSKAP nocodazole-treated Rpe1-GFP cells (2 μ M nocodazole, 3 h) stained for CREST (yellow), chromosomes (purple), and tubulin (red). Scale bar, 10 μ m.

(E) K-K distance for individual sister pairs in the control and siSKAP cells treated with nocodazole (Mann-Whitney test) (n = number of kinetochore pairs).

(F and G) Laser ablation (yellow X) of k-fiber near a kinetochore releases tension, if present, across a sister pair (schematic cartoon) (F), as shown in representative kymograph images of K-K distance relaxation upon k-fiber ablation (yellow arrowhead) in Rpe1-GFP cells (G). Scale bar, 10 μ m (G).

(H and I) K-K distance relaxation (decrease) post-ablation as a function of K-K distance pre-ablation (linear regression lines for each condition) (H) or as a direct comparison (I) (Student's t test) in the control versus siSKAP cells (n = number of ablations, one ablation per cell).

See also Figure S2.




Figure 3. SKAP decreases kinetochore friction on polymerizing microtubules

(A) k-fiber ablation assay to isolate kinetochores associated with polymerizing microtubules: laser ablation (yellow X) leads to transient retraction of the kinetochore pair, the creation of new minus-ends leads to the recruitment of dynein (light blue), which pulls on a kinetochore and leads the microtubules attached to its sister to polymerize (pink).

(legend continued on next page)

K-K distance was a specific effect of SKAP silencing rather than an off-target effect of the siRNA used in this study (Figure S2A).

In principle, an increase in K-K distance could stem from increased tension at the kinetochore-microtubule interface or decreased centromere stiffness.^{7,44} To test whether siSKAP cells could arrest at mitosis and thereby indirectly undergo cohesion fatigue and centromere softening,⁴⁴ we imaged and measured mitotic duration (nuclear envelope breakdown [NEB] to anaphase onset) in Rpe1-GFP cells. The mitotic duration was ~24 min in the control and ~8 min longer in siSKAP cells (23.9 ± 5.8 versus 31.4 ± 15.0 min; Figure 2C). However, when we artificially induced mitotic arrest using the proteasome inhibitor MG132 (10 μM), the K-K distance only detectably increased starting from 106 min post MG132 (Figures S2B and S2C).⁴⁴ Considering that just 6% of siSKAP cells had a mitotic duration longer than 100 min (Figure 2C), cohesion fatigue is unlikely responsible for increasing the K-K distance in siSKAP cells, as the mitotic delay observed in siSKAP cells is not long enough to induce this effect. To test whether SKAP could more directly affect centromere stiffness, we compared the chromosome movements in these MG132-treated cells with those in siSKAP cells (Figure S2). MG132-treated cells selected for their high mean K-K distance (Figure S2B), indicative of low centromere stiffness, had poor sister kinetochore velocity correlation, indistinguishable from siSKAP cells, but kinetochore velocity indistinguishable from the control (Figures S2D and S2E). Thus, although an effect on centromere stiffness cannot be fully excluded, SKAP cannot simply increase centromere stiffness. Consistent with SKAP not affecting centromere stiffness, treating Rpe1-GFP cells with nocodazole to remove microtubules and spindle forces led to reduced K-K distances that were indistinguishable between siSKAP and the control (0.7 ± 0.2 μm siSKAP versus 0.7 ± 0.1 μm control; Figures 2D and 2E). Thus, the increased K-K distance in siSKAP kinetochores depends on microtubules. Supporting this idea, laser ablating k-fibers in Rpe1-GFP cells to release spindle forces (Figures 2F and 2G) indicates that siSKAP kinetochores are under more tension than the control. As expected for a spring, sister pairs with higher K-K distances pre-ablation relaxed more post-ablation than those with lower K-K distances in both control and siSKAP cells (Figure 2H). Notably, K-K distance relaxed more post-ablation in siSKAP kinetochores than in the control (0.5 ± 0.2 μm versus 0.4 ± 0.1 μm; Figure 2I). This difference indicates that siSKAP kinetochores are under higher tension at the

kinetochore-microtubule interface and is consistent with sister kinetochores coordinating movement more poorly (Figures 1E–1G). Thus, SKAP reduces tension at the kinetochore-microtubule interface, and not all microtubule couplers at the kinetochore increase tension at this interface.

SKAP decreases kinetochore friction on polymerizing microtubules

To decrease tension at the kinetochore-microtubule interface, SKAP could either reduce passive, frictional force; active, energy-consuming force generated at this interface; or both. At metaphase, the back sister kinetochore is typically bound to polymerizing microtubules through a largely passive interface and the front sister to depolymerizing microtubules through an interface that is both active and passive.⁶ Thus, decoupling the roles of SKAP at passive versus active interfaces requires uncoupling the effect of SKAP depletion at kinetochores in polymerizing and depolymerizing microtubules.

Since both sisters are attached together, exerting force on each other and holding on to microtubules in opposite polymerization states, we turned to laser ablation to decouple their responses. To probe kinetochores bound to polymerizing microtubules and generating passive force, we ablated k-fibers to trigger spindle-based force generation on a sister pair.¹⁸ k-fiber ablation generates new microtubule minus-ends that are recognized by dynein and pulled to the spindle pole (Figure 3A), which exerts an “external” force on the attached sister pair and induces microtubule polymerization at the back (away from the pole) kinetochore.^{45,46} Assuming that this dynein pulling force is comparable between conditions, the velocity of this back kinetochore reports on the passive, frictional force at its microtubule interface.¹⁸ We ablated k-fibers in Rpe1-GFP cells with and without SKAP (Figure 3B; Videos S3 and S4) and tracked both sisters. The velocity of the front kinetochore was indistinguishable in siSKAP and control cells (3.2 ± 1.5 μm/min siSKAP versus 3.0 ± 1.4 μm/min in controls; Figures 3C and 3E). This is consistent with the dynein-generated force not being affected in siSKAP cells. By contrast, the back kinetochore moved slower in siSKAP cells than in the control (2.0 ± 0.9 versus 2.8 ± 1.2 μm/min; Figures 3D and 3E). Although the front and back kinetochores moved at indistinguishable velocities in the control cells (3.0 ± 1.4 μm/min in front versus 2.8 ± 1.2 μm/min in back kinetochores), the back kinetochores moved at lower velocities than the front ones in siSKAP cells (2.0 ± 0.9 versus 3.2 ± 1.5 μm/min) (Figure 3E). This led to a

(B) Representative time-lapse of k-fiber ablation in Rpe1-GFP control and siSKAP cells. The white lines mark the kinetochore positions pre-ablation at the top, the red lines represent the movement of sister kinetochores. Scale bar, 10 μm.

(C) Individual tracks from control and siSKAP front kinetochores moving post-ablation, showing kinetochore displacement from its switching position after dynein pulling (time = 0 corresponds to the time point of the switch to poleward movement), with linear regression fits (straight lines) (ANCOVA).

(D) Individual tracks from control and siSKAP back kinetochores moving post-ablation, showing kinetochore displacement from its switching position after dynein pulling (time = 0 corresponds to the time point of the switch to away from the pole movement), with linear regression fits (straight lines) (ANCOVA).

(E) Front (lined boxes) and back (clear boxes) kinetochore speed distribution post-ablation in control and siSKAP kinetochores (Student's t test) from data in (C) and (D).

(F) K-K distance at the time the back kinetochore switched to away-from-pole movement post-ablation in control and siSKAP kinetochore pairs (Mann-Whitney test).

In (C)–(F), n = number of ablations, one ablation per cell.

(G) Representative immunofluorescence images of Rpe1-GFP control and siSKAP cells stained for Hec1-S69 phosphorylation (red), tubulin (blue), and Hec1 (yellow). Scale bar, 5 μm.

(H) p-Hec1-S69 kinetochore intensity relative to Hec1 kinetochore intensity in control and siSKAP cells (Student's t test; n = number of cells).

See also Figure S3 and Videos S3 and S4.

persistent increase in the K-K distance during this sister kinetochore movement in siSKAP kinetochore pairs (Figure S3). Notably, the K-K distance at the time of back kinetochore directional switching (starting to move away from the pole) was higher in siSKAP kinetochores than in the control ($1.4 \pm 0.5 \mu\text{m}$ versus $1.0 \pm 0.2 \mu\text{m}$; Figure 3F), again suggesting that siSKAP kinetochores are less sensitive to force changes. Together, these findings indicate that SKAP decreases the friction between kinetochores and polymerizing microtubules.

Hec1 dephosphorylation is known to increase friction at the attachment interface.^{17,18} Hec1-S69 phosphorylation (by Aurora A) is maintained during mitosis and is required and sufficient to preserve kinetochore movement dynamics at metaphase.²⁵ Therefore, we tested whether SKAP depletion affected the level of Hec1-S69 phosphorylation. Immunofluorescence quantification showed that SKAP depletion did not affect the kinetochore levels of Hec1-S69 phosphorylation (Figures 3G and 3H), indicating that the friction regulation by SKAP occurs independently of Hec1-S69 phosphorylation. Together, these findings indicate that SKAP, in contrast to the proposed role of other microtubule-binding proteins at the kinetochore,^{17,18,20,28} decreases the friction between kinetochores and polymerizing microtubules and does so through a mechanism independent of that setting Hec1's basal dynamic state.²⁵ SKAP could either do so by increasing the microtubule tip polymerization dynamics, producing an apparent change in friction, or by directly reducing the kinetochore friction on the microtubule lattice.

SKAP increases k-fiber depolymerization velocity and kinetochore force responsiveness

Given that SKAP regulates the kinetochore's frictional interface with polymerizing microtubules (Figure 3), we asked whether it also regulates its interface with depolymerizing ones. To decouple both attached sister kinetochores and isolate SKAP's role in depolymerizing microtubules, we laser ablated one kinetochore in a sister pair in Rpe1-GFP cells and tracked the movement of the remaining sister. As expected, the remaining sister was pulled poleward by depolymerizing microtubules and reversed direction near the pole, pushed by polar ejection forces^{3,9,12} (Figures 4A and 4B; Video S5). The poleward velocity of siSKAP kinetochores was slower than that in the control (2.1 ± 0.6 versus $3.5 \pm 0.8 \mu\text{m}/\text{min}$; Figures 4B–4D; Videos S5 and S6). This indicates that SKAP increases the velocity of microtubule depolymerization at the kinetochore interface. Here too, it could do so either by lowering the friction on the microtubule lattice or by increasing the depolymerization dynamics and thereby decreasing the apparent friction. Consistent with SKAP acting directly at the interface, SKAP depletion did not detectably change the kinetochore levels of key microtubule plus-end depolymerases involved in metaphase kinetochore movements, MCAK and Kif18A^{47,48} (Figure S4). Further, in the same kinetochore ablation experiments, siSKAP kinetochores switched direction (to away from pole movement) closer to the pole than the control kinetochores as they got pushed by polar ejection forces (3.2 ± 1.2 versus $4.0 \pm 1.8 \mu\text{m}$; Figure 4E). Thus, SKAP increases the kinetochore's force sensitivity when it is bound to depolymerizing microtubules, favoring a switch to the polymerization state, as observed after k-fiber ablation (Figure 3F) and during metaphase oscillations (Figure 1G). Together, these

findings indicate that Astrin-SKAP does not increase the kinetochore's grip on microtubules as other microtubule-binding proteins are thought to do but instead reduces grip, lowering friction and effectively lubricating the interface, making it more dynamic and force responsive.

DISCUSSION

Faithful chromosome segregation requires kinetochores to hold on to dynamic microtubules as they grow and shrink. Here, we ask the following: how does the mammalian kinetochore-microtubule interface stay dynamic and force responsive while maintaining a strong grip on microtubules? Mechanisms increasing the grip as attachments mature are being actively studied, including Ndc80 dephosphorylation^{17,18} and Ska recruitment.^{19,20} Here, we combine molecular and mechanical perturbations to define the mechanical role of SKAP at this interface, and we show that it, in turn, decreases the grip at the interface. We demonstrate that SKAP increases sister kinetochore mobility and coordination in metaphase (Figure 1) and decreases the tension at the kinetochore-microtubule interface (Figure 2). We show that SKAP increases the velocity at which kinetochores move on both polymerizing (Figure 3) and depolymerizing (Figure 4) microtubules and that it makes the attachment more responsive to force changes (Figures 3 and 4). Together, our findings indicate that SKAP reduces the friction at the kinetochore-microtubule interface, effectively lubricating it. As such, and given SKAP's arrival as kinetochores biorient,^{29,30} we propose that SKAP keeps the correct kinetochore-microtubule attachments dynamic to preserve them as they stabilize and mature.^{17,19} The association of SKAP mutations and changes in expression with some cancers and aneuploidy^{38,39,49} is consistent with SKAP's lubrication function being key for faithful chromosome segregation.

Our work indicates that SKAP lowers the friction at the kinetochore-microtubule interface, raising the question of what function a lower friction could serve. Basal levels of attachment dynamics are essential for accurate chromosome segregation.^{25,50} Ska recruitment and the gradual dephosphorylation of Hec1 increases the kinetochore's grip on microtubules during mitosis.^{17–20} In contrast to other Hec1 phosphosites, S69 is persistently phosphorylated during mitosis, maintaining basal dynamics.²⁵ Yet these dynamics are lost and friction increases upon SKAP depletion (Figures 1, 3, and 4), although the phosphorylation levels of Hec1-S69 are not affected by this depletion (Figures 3G and 3H). Thus, SKAP acts as a lubricant independently of Hec1-S69 phosphorylation. By reducing friction, SKAP may increase microtubule dynamics, increasing microtubule depolymerization (Figures 4B–4D) and poleward flux.³³ To our knowledge, SKAP is the first microtubule binder proposed to decrease the friction at the kinetochore-microtubule interface. We propose that using SKAP to lower the friction, instead of loosening Ndc80-Ska's grip to lower the friction, allows the cell to keep a dynamic kinetochore-microtubule interface without losing the grip of microtubules and attachment stability, by having one molecule specialized for each activity. Indeed, having distinct mechanisms to tune grip and dynamics may lead to finer regulatory control “knobs” to preserve stable attachments.




Figure 4. SKAP increases k-fiber depolymerization velocity and kinetochore's force responsiveness

(A) Kinetochore ablation assay to isolate kinetochores associated with depolymerizing microtubules: laser ablation (yellow X) of one sister leads to the other sister moving poleward as its microtubules depolymerize (purple) and to later move away from the pole as polar ejection forces (green arrowheads) increase and microtubules rescue and polymerize (pink).

(B) Representative time-lapse images of kinetochore ablation (yellow arrow) in the control (top) and siSKAP (bottom) Rpe1-GFP cells, with remaining sister kinetochore (white arrow) and centrioles (red arrows) marked and kinetochore directional switch marked (purple arrow). Scale bar, 10 µm.

(C) Distance to the position at ablation as a function of time for individual kinetochores post-ablation ($t = 0$ corresponds to the first time point post-ablation) in the control and siSKAP cells, with linear regression fits (straight lines) (ANCOVA).

(D) The average speed of individual kinetochores post sister ablation in the control and siSKAP cells (Mann-Whitney test).

(E) Kinetochore distance from the spindle pole of individual control and siSKAP cells at the time of direction switch from poleward to away-from-pole movement (rescue) post-ablation (Student's *t* test), with a smaller distance typically reflecting a higher force at rescue.

In (C)–(E), n = number of ablations, one ablation per cell.

See also [Figure S4](#) and [Videos S5](#) and [S6](#).

By lubricating the correct, bioriented attachments, SKAP could in principle help preserve them and prevent their increased microtubule affinity from making them less mobile and less responsive to force. The presence of more dynamic, lower friction interactions with microtubules could help stabilize attachments under force by increasing their adaptability to force changes. The force responsiveness of attachments is essential for accurate chromosome movement, alignment, and sister kinetochore coordination,^{7,12,51} and SKAP depleted cells have

chromosome alignment and segregation defects.^{29–31} Indeed, we show that SKAP makes kinetochores more mobile ([Figures 1C](#) and [1D](#)) and the kinetochore-microtubule interface more sensitive to force changes ([Figures 1E–1G](#), [3F](#), and [4E](#)). This could explain why siSKAP attachments are under higher tension ([Figure 2](#)) and inefficient at dissipating force ([Figures 3B–3E](#)) and at responding to sister movement ([Figures 1E–1G](#) and [3F](#)), and it suggests that a function of lowering friction could be to make kinetochores responsive to




Figure 5. Models for the Astrin-SKAP complex decreasing friction at the mammalian kinetochore-microtubule interface

Representation of the kinetochore-microtubule interface in the absence (left) and presence (right) of SKAP. Two models for how Astrin-SKAP (yellow) could increase the friction at the kinetochore-microtubule interface (right). In model A (top right), SKAP affects how Ndc80 (black) or Ndc80-Ska (purple) complexes bind microtubules (dashed black and purple), decreasing their friction on microtubules and indirectly reducing the attachment friction and increasing dynamics. In model B (bottom right), SKAP directly binds microtubules, with Ndc80-SKAP and Ndc80-Ska competing for microtubule binding with similar affinities (binding energy) but with Ndc80-SKAP moving on microtubules with lower friction (lower transition state energy in moving between lattice binding sites). In both models, the more SKAP molecules are at the kinetochore-microtubule interface and the higher the sensitivity to force (center).

force. While additional work will be needed to reveal how SKAP increases force responsiveness, in principle, lowering friction would be sufficient to do so.

Although SKAP reduces the friction at the kinetochore-microtubule interface (Figures 3A–3E), the mechanism by which it does so is not clear. SKAP depletion could in principle increase friction by reducing microtubule dynamics. However, lowering microtubule dynamics globally with drugs (such as taxol or eribulin) decreases interkinetochore tension (or K-K distance) rather than increasing it as we see for siSKAP (Figure 2).^{52,53} Thus, decreasing microtubule dynamics is not sufficient to recapitulate the effects of SKAP depletion. Further, we could not detect any change in the kinetochore recruitment of key microtubule dynamics regulators (MCAK and Kif18A) in siSKAP cells (Figure S4), and changing their activity is not consistent with our findings: MCAK depletion decreases both interkinetochore tension and microtubule depolymerization,⁴⁷ and Kif18A depletion increases kinetochore velocity and oscillation amplitudes and decreases switching rates,^{48,51} and we do not see these observables change as such (Figures 1C, 1D, 2, 3F, and 4C–4E). Alternatively, SKAP might affect microtubule dynamics through its described interaction with the plus-end factor EB1.^{33,54} Contradicting this possibility, SKAP's interaction with EB1 is not needed for SKAP's role in chromosome alignment and spindle assembly, suggesting that SKAP's role at the kinetochore-microtubule interface is independent of its interaction with EB1.⁵⁵ Thus, although we cannot exclude the possibility that SKAP regulates the kinetochore activity of these or other microtubule dynamics regulators or directly affects microtubule dynamics, a change in microtubule dynamics alone is not sufficient to recapitulate all the behaviors of siSKAP cells. Together, these findings are consistent with SKAP playing a mechanical role at the kinetochore-microtubule interface and not simply regulating microtubule dynamics.

Mechanistically, SKAP could regulate the kinetochore-microtubule friction in different ways. One way in which it could do so is by changing the sliding friction of the other kinetochore proteins on microtubules. SKAP could induce conformational changes in Ndc80 or Ska complexes, which lower their friction on

microtubules (model A; Figure 5), independently of Hec1-S69 phosphoregulation (Figures 3G and 3H). Alternatively, SKAP could directly bind microtubules and act as a kinetochore-microtubule coupler, as suggested by *in vitro* work.^{30,34,35} Accordingly, the microtubule-binding domain of SKAP is essential for its role in chromosome alignment and segregation.⁵⁵ As such, SKAP could then compete out other microtubule binders that have a higher friction on microtubules (model B; Figure 5), as microtubule affinity (binding energy) and friction coefficient (transition state energy in moving between lattice binding sites) are not strictly coupled.⁵⁶ The affinity of Ndc80-SKAP on microtubules³⁵ is similar, although lower than that of Ndc80-Ska,¹⁶ suggesting that both complexes could compete for microtubule binding. If Ndc80-SKAP had a lower friction on microtubules than Ndc80-Ska, SKAP kinetochore localization could lower the friction at the kinetochore-microtubule interface. In principle, the above models hold independently of how SKAP is recruited to the kinetochore, for example, through Ndc80^{30,35} or Mis13.³³ Whether SKAP lowers other kinetochore proteins' (Ndc80, Ska) friction on microtubules (model A) and whether SKAP and other kinetochore proteins compete for microtubule binding and generate different friction (model B) are non-exclusive models and a rich area of future study.

Across scales in biology, diverse interfaces need to be both strong and dynamic. For example, motor proteins must walk on their tracks without letting them go, and cell-cell junctions must robustly persist yet remodel. Understanding the mechanics of such interfaces will require understanding not only how strong, robust interactions are achieved but also how the interface can remain dynamic. The latter may require looking beyond mechanisms that grip, as is the case for SKAP, which makes the human kinetochore-microtubule interface more dynamic.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE

● RESOURCE AVAILABILITY

- Lead contact
- Materials availability
- Data and code availability

● EXPERIMENTAL MODEL AND SUBJECT DETAILS

● METHOD DETAILS

- Cell culture, siRNA transfection and drug treatments in Rpe1 cells
- Immunofluorescence and immunoblotting
- Microscopy and laser ablation
- Study design and data inclusion criteria

● QUANTIFICATION AND STATISTICAL ANALYSIS

- Analyzing fluorescence intensity
- Analyzing kinetochore behaviors
- Statistical analysis
- Video preparation

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.cub.2022.04.061>.

ACKNOWLEDGMENTS

We thank Iain Cheeseman for the anti-SKAP antibody and discussion, Jennifer DeLuca for the Hec1-pS69 antibody, Alexey Khodjakov for the Rpe1-GFP (GFP-CenpA and centrin1-GFP) cells, and Charles Asbury and Rob Phillips for discussions. We thank the members of the Dumont Lab for their discussions and critical reading of the manuscript. This work was supported by NIH DP2GM119177, NIH R01GM134132, NIH R35GM136420, NSF CAREER 1554139, NSF 1548297 Center for Cellular Construction, the Rita Allen Foundation, and the Chan Zuckerberg Biohub (S.D.).

AUTHOR CONTRIBUTIONS

Conceptualization, M.R.-S. and S.D.; methodology, M.R.-S.; software, M.R.-S. and P.S.; validation, M.R.-S.; formal analysis, M.R.-S.; investigation, M.R.-S. and R.S. (Figures 2C, 3G, and S4); resources, S.D.; data curation, M.R.-S.; writing – original draft, M.R.-S.; writing – review & editing, M.R.-S. and S.D.; visualization, M.R.-S.; supervision, M.R.-S. and S.D.; funding acquisition, S.D.

DECLARATION OF INTERESTS

The authors declare no conflict of interest.

Received: December 16, 2021

Revised: March 17, 2022

Accepted: April 21, 2022

Published: May 16, 2022

REFERENCES

1. Wendell, K.L., Wilson, L., and Jordan, M.A. (1993). Mitotic block in HeLa cells by vinblastine: ultrastructural changes in kinetochore-microtubule attachment and in centrosomes. *J. Cell Sci.* 104, 261–274.
2. McEwen, B.F., Heagle, A.B., Cassels, G.O., Buttle, K.F., and Rieder, C.L. (1997). Kinetochore fiber maturation in PtK1 cells and its implications for the mechanisms of chromosome congression and anaphase onset. *J. Cell Biol.* 137, 1567–1580.
3. Khodjakov, A., and Rieder, C.L. (1996). Kinetochores moving away from their associated pole do not exert a significant pushing force on the chromosome. *J. Cell Biol.* 135, 315–327.
4. Grishchuk, E.L., and McIntosh, J.R. (2006). Microtubule depolymerization can drive poleward chromosome motion in fission yeast. *EMBO J.* 25, 4888–4896.
5. Joglekar, A.P., Bloom, K.S., and Salmon, E.D. (2010). Mechanisms of force generation by end-on kinetochore-microtubule attachments. *Curr. Opin. Cell Biol.* 22, 57–67.
6. Dumont, S., Salmon, E.D., and Mitchison, T.J. (2012). Deformations within moving kinetochores reveal different sites of active and passive force generation. *Science* 337, 355–358.
7. Wan, X., Cimini, D., Cameron, L.A., and Salmon, E.D. (2012). The coupling between sister kinetochore directional instability and oscillations in centromere stretch in metaphase PtK1 cells. *Mol. Biol. Cell* 23, 1035–1046.
8. Rago, F., and Cheeseman, I.M. (2013). Review series: the functions and consequences of force at kinetochores. *J. Cell Biol.* 200, 557–565.
9. McNeill, P.A., and Berns, M.W. (1981). Chromosome behavior after laser microirradiation of a single kinetochore in mitotic PtK2 cells. *J. Cell Biol.* 88, 543–553.
10. Skibbens, R.V., Rieder, C.L., and Salmon, E.D. (1995). Kinetochore motility after severing between sister centromeres using laser microsurgery: evidence that kinetochore directional instability and position is regulated by tension. *J. Cell Sci.* 108, 2537–2548.
11. Rieder, C.L., and Salmon, E.D. (1994). Motile kinetochores and polar ejection forces dictate chromosome position on the vertebrate mitotic spindle. *J. Cell Biol.* 124, 223–233.
12. Ke, K., Cheng, J., and Hunt, A.J. (2009). The distribution of polar ejection forces determines the amplitude of chromosome directional instability. *Curr. Biol.* 19, 807–815.
13. Bakhroum, S.F., and Compton, D.A. (2012). Kinetochores and disease: keeping microtubule dynamics in check. *Curr. Opin. Cell Biol.* 24, 64–70.
14. DeLuca, J.G., Dong, Y., Hergert, P., Strauss, J., Hickey, J.M., Salmon, E.D., and McEwen, B.F. (2005). Hec1 and Nuf2 are core components of the kinetochore outer plate essential for organizing microtubule attachment sites. *Mol. Biol. Cell* 16, 519–531.
15. Cheeseman, I.M., and Desai, A. (2008). Molecular architecture of the kinetochore-microtubule interface. *Nat. Rev. Mol. Cell Biol.* 9, 33–46.
16. Schmidt, J.C., Arthanari, H., Boeszoermenyi, A., Dashkevich, N.M., Wilson-Kubalek, E.M., Monnier, N., Markus, M., Oberer, M., Milligan, R.A., Bathe, M., et al. (2012). The kinetochore-bound Ska1 complex tracks depolymerizing microtubules and binds to curved protofilaments. *Dev. Cell* 23, 968–980.
17. Zaytsev, A.V., Sundin, L.J.R., DeLuca, K.F., Grishchuk, E.L., and DeLuca, J.G. (2014). Accurate phosphoregulation of kinetochore-microtubule affinity requires unconstrained molecular interactions. *J. Cell Biol.* 206, 45–59.
18. Long, A.F., Udy, D.B., and Dumont, S. (2017). Hec1 tail phosphorylation differentially regulates mammalian kinetochore coupling to polymerizing and depolymerizing microtubules. *Curr. Biol.* 27, 1692–1699.e3.
19. Auckland, P., Clarke, N.I., Royle, S.J., and McAinsh, A.D. (2017). Congressing kinetochores progressively load Ska complexes to prevent force-dependent detachment. *J. Cell Biol.* 216, 1623–1639.
20. Huis In 't Veld, P.J., Volkov, V.A., Stender, I.D., Musacchio, A., Dogterom, M., Huis, P.J., Volkov, V.A., Stender, I.D., Musacchio, A., and Dogterom, M. (2019). Molecular determinants of the Ska-Ndc80 interaction and their influence on microtubule tracking and force-coupling. *eLife* 8, e49539.
21. Long, A.F., Kuhn, J., and Dumont, S. (2019). The mammalian kinetochore-microtubule interface: robust mechanics and computation with many microtubules. *Curr. Opin. Cell Biol.* 60, 60–67.
22. Cheeseman, I.M., Chappie, J.S., Wilson-Kubalek, E.M., and Desai, A. (2006). The conserved KMN network constitutes the core microtubule-binding site of the kinetochore. *Cell* 127, 983–997.
23. Powers, A.F., Franck, A.D., Gestaut, D.R., Cooper, J., Graczyk, B., Wei, R.R., Wordeman, L., Davis, T.N., and Asbury, C.L. (2009). The Ndc80 kinetochore complex forms load-bearing attachments to dynamic microtubule tips via biased diffusion. *Cell* 136, 865–875.

24. Zaytsev, A.V., Mick, J.E., Maslennikov, E., Nikashin, B., DeLuca, J.G., and Grishchuk, E.L. (2015). Multisite phosphorylation of the NDC80 complex gradually tunes its microtubule-binding affinity. *Mol. Biol. Cell* **26**, 1829–1844.
25. DeLuca, K.F., Meppelink, A., Broad, A.J., Mick, J.E., Peersen, O.B., Pektas, S., Lens, S.M.A., and DeLuca, J.G. (2018). Aurora A kinase phosphorylates Hec1 to regulate metaphase kinetochore-microtubule dynamics. *J. Cell Biol.* **217**, 163–177.
26. Hanisch, A., Silljé, H.H., and Nigg, E.A. (2006). Timely anaphase onset requires a novel spindle and kinetochore complex comprising Ska1 and Ska2. *EMBO J.* **25**, 5504–5515.
27. Welburn, J.P.I., Grishchuk, E.L., Backer, C.B., Wilson-Kubalek, E.M., Yates, J.R., and Cheeseman, I.M. (2009). The human kinetochore Ska1 complex facilitates microtubule depolymerization-coupled motility. *Dev. Cell* **16**, 374–385.
28. Helgeson, L.A., Zelter, A., Riffle, M., MacCoss, M.J., Asbury, C.L., and Davis, T.N. (2018). Human Ska complex and Ndc80 complex interact to form a load-bearing assembly that strengthens kinetochore-microtubule attachments. *Proc. Natl. Acad. Sci. USA* **115**, 2740–2745.
29. Fang, L., Seki, A., and Fang, G. (2009). SKAP associates with kinetochores and promotes the metaphase-to-anaphase transition. *Cell Cycle* **8**, 2819–2827.
30. Schmidt, J.C., Kiyomitsu, T., Hori, T., Backer, C.B., Fukagawa, T., and Cheeseman, I.M. (2010). Aurora B kinase controls the targeting of the Astrin-SKAP complex to bioriented kinetochores. *J. Cell Biol.* **191**, 269–280.
31. Dunsch, A.K., Linnane, E., Barr, F.A., and Gruneberg, U. (2011). The astrin-kinastrin/SKAP complex localizes to microtubule plus ends and facilitates chromosome alignment. *J. Cell Biol.* **192**, 959–968.
32. Cimini, D., Wan, X., Hirel, C.B., and Salmon, E.D. (2006). Aurora kinase promotes turnover of kinetochore microtubules to reduce chromosome segregation errors. *Curr. Biol.* **16**, 1711–1718.
33. Wang, X., Zhuang, X., Cao, D., Chu, Y., Yao, P., Liu, W., Liu, L., Adams, G., Fang, G., Dou, Z., et al. (2012). Mitotic regulator SKAP forms a link between kinetochore core complex KMN and dynamic spindle microtubules. *J. Biol. Chem.* **287**, 39380–39390.
34. Friese, A., Faesen, A.C., Huis In 't Veld, P.J., Fischböck, J., Prumbaum, D., Petrovic, A., Raunser, S., Herzog, F., and Musacchio, A. (2016). Molecular requirements for the inter-subunit interaction and kinetochore recruitment of SKAP and Astrin. *Nat. Commun.* **7**, 11407.
35. Kern, D.M., Monda, J.K., Su, K.-C., Wilson-Kubalek, E.M., and Cheeseman, I.M. (2017). Astrin-SKAP complex reconstitution reveals its kinetochore interaction with microtubule-bound Ndc80. *eLife* **6**, e26866.
36. Gao, H., Pan, Q., Wang, Y., and Chen, Q. (2021). Impact of KMN network genes on progression and prognosis of non-small cell lung cancer. *Anticancer Drugs* **33**, e398–e408.
37. Yu, D.C., Chen, X.Y., Li, X., Zhou, H.Y., Yu, D.Q., Yu, X.L., Hu, Y.C., Zhang, R.H., Zhang, X.B., Zhang, K., et al. (2021). Transcript levels of spindle and kinetochore-associated complex 1/3 as prognostic biomarkers correlated with immune infiltrates in hepatocellular carcinoma. *Sci. Rep.* **11**, 11165.
38. Lee, C.S., Bhaduri, A., Mah, A., Johnson, W.L., Ungewickell, A., Aros, C.J., Nguyen, C.B., Rios, E.J., Siphrašvili, Z., Straight, A., et al. (2014). Recurrent point mutations in the kinetochore gene KNSTRN in cutaneous squamous cell carcinoma. *Nat. Genet.* **46**, 1060–1062.
39. Deng, P., Zhou, R., Zhang, J., and Cao, L. (2021). Increased expression of KNSTRN in lung adenocarcinoma predicts poor prognosis: a bioinformatics analysis based on TCGA data. *J. Cancer* **12**, 3239–3248.
40. Magidson, V., O'Connell, C.B., Lončarek, J., Paul, R., Mogilner, A., and Khodjakov, A. (2011). The spatial arrangement of chromosomes during prometaphase facilitates spindle assembly. *Cell* **146**, 555–567.
41. DeLuca, J.G., Gall, W.E., Ciferri, C., Cimini, D., Musacchio, A., and Salmon, E.D. (2006). Kinetochore microtubule dynamics and attachment stability are regulated by Hec1. *Cell* **127**, 969–982.
42. Raaijmakers, J.A., Tanenbaum, M.E., Maia, A.F., and Medema, R.H. (2009). RAMA1 is a novel kinetochore protein involved in kinetochore-microtubule attachment. *J. Cell Sci.* **122**, 2436–2445.
43. Huang, Y., Wang, W., Yao, P., Wang, X., Liu, X., Zhuang, X., Yan, F., Zhou, J., Du, J., Ward, T., et al. (2012). CENP-E kinesin interacts with SKAP protein to orchestrate accurate chromosome segregation in mitosis. *J. Biol. Chem.* **287**, 1500–1509.
44. Daum, J.R., Potapova, T.A., Sivakumar, S., Daniel, J.J., Flynn, J.N., Rankin, S., and Gorbisky, G.J. (2011). Cohesion fatigue induces chromatid separation in cells delayed at metaphase. *Curr. Biol.* **21**, 1018–1024.
45. Sikirzhyski, V., Magidson, V., Steinman, J.B., He, J., Le Berre, M., Tikhonenko, I., Ault, J.G., McEwen, B.F., Chen, J.K., Sui, H., et al. (2014). Direct kinetochore-spindle pole connections are not required for chromosome segregation. *J. Cell Biol.* **206**, 231–243.
46. Elting, M.W., Hueschen, C.L., Udy, D.B., and Dumont, S. (2014). Force on spindle microtubule minus ends moves chromosomes. *J. Cell Biol.* **206**, 245–256.
47. Wordeman, L., Wagenbach, M., and Dassow, G. Von. (2007). MCAK facilitates chromosome movement by promoting kinetochore microtubule turnover. *J. Cell Biol.* **179**, 869–879.
48. Stumpff, J., Von Dassow, G., Wagenbach, M., Asbury, C., and Wordeman, L. (2008). The Kinesin-8 motor Kif18A suppresses kinetochore movements to control mitotic chromosome alignment. *Dev. Cell* **14**, 252–262.
49. Jaju, P.D., Nguyen, C.B., Mah, A.M., Atwood, S.X., Li, J., Zia, A., Chang, A.L.S., Oro, A.E., Tang, J.Y., Lee, C.S., et al. (2015). Mutations in the kinetochore gene KNSTRN in basal cell carcinoma. *J. Invest. Dermatol.* **135**, 3197–3200.
50. Bakhom, S.F., Thompson, S.L., Manning, A.L., and Compton, D.A. (2009). Genome stability is ensured by temporal control of kinetochore-microtubule dynamics. *Nat. Cell Biol.* **11**, 27–35.
51. Stumpff, J., Wagenbach, M., Franck, A., Asbury, C.L., and Wordeman, L. (2012). Kif18A and Chromokinesins confine centromere movements via microtubule growth suppression and spatial control of kinetochore tension. *Dev. Cell* **22**, 1017–1029.
52. Waters, J.C., Chen, R.H., Murray, A.W., and Salmon, E.D. (1998). Localization of Mad2 to kinetochores depends on microtubule attachment, not tension. *J. Cell Biol.* **141**, 1181–1191.
53. Okouneva, T., Azarenko, O., Wilson, L., Littlefield, B.A., and Jordan, M.A. (2008). Inhibition of centromere dynamics by eribulin (E7389) during mitotic metaphase. *Mol. Cancer Ther.* **7**, 2003–2011.
54. Tamura, N., Simon, J.E., Nayak, A., Shenoy, R.T., Hiroi, N., Boillot, V., Funahashi, A., and Draviam, V.M. (2015). A proteomic study of mitotic phase-specific interactors of EB1 reveals a role for SXIP-mediated protein interactions in anaphase onset. *Biol. Open* **4**, 155–169.
55. Kern, D.M., Nicholls, P.K., Page, D.C., and Cheeseman, I.M. (2016). A mitotic SKAP isoform regulates spindle positioning at astral microtubule plus ends. *J. Cell Biol.* **213**, 315–328.
56. Forth, S., Hsia, K.C., Shimamoto, Y., and Kapoor, T.M. (2014). Asymmetric friction of nonmotor MAPs can lead to their directional motion in active microtubule networks. *Cell* **157**, 420–432.
57. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682.
58. Meijering, E., Dzyubachyk, O., and Smal, I. (2012). *Methods for Cell and Particle Tracking*, First Edition (Elsevier).

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit anti-SKAP	Origene	TA333584
Rabbit anti-SKAP	I. Cheeseman lab	Kern et al. ⁵⁵
Mouse anti- α -tubulin (DM1A)	Sigma-Aldrich	T6199; RRID: AB_477583
Rabbit anti-Kif18A	Bethyl	A301-080A-M; RRID: AB_2296551
Rabbit anti-MCAK	cytoskeleton	AKIN05; RRID: AB_10708227
Rabbit anti-p-Hec1-S69	J. DeLuca lab	DeLuca et al. ²⁵
Human anti-CREST	Antibodies Incorporated	15-234-0001; RRID: AB_2687472
Mouse anti-Hec1	Abcam	ab3613; RRID: AB_303949
Rat anti- α -tubulin	Bio-rad	MCA77G; RRID: AB_325003
Goat anti-mouse IgG Alexa Fluor 405, 488 and 568	Invitrogen	A31553, A11001 and A11004
Goat anti-rabbit IgG Alexa Fluor 405 and 568	Invitrogen	A31556 and A11011
Goat anti-rat IgG Alexa Fluor 488	Invitrogen	A11006
Goat anti-human IgG Alexa Fluor 568	Invitrogen	A21090
Chemicals, peptides, and recombinant proteins		
Nocodazole	Sigma-Aldrich	M1404-50MG
MG132	EMB Millipore	474790-5MG
SiR-tubulin	Cytoskeleton	CY-SC002
Experimental models: Cell lines		
Rpe1-GFP cells	A. Khodjakov lab	Magidson et al. ⁴⁰
Oligonucleotides		
siRNA SKAP: 5'-AGGCUACAAACCACUGAGUAA-3'	Sigma-Aldrich	Dunsch et al. ³¹
siRNA SKAP2: 5'-GAAAGAGUCCGAUUCUAG-3'	Sigma-Aldrich	Schmidt et al. ³⁰
siRNA luciferase: 5' CGUACGCGGAUACUUCGA 3'	Sigma-Aldrich	WD01818022
Software and algorithms		
FIJI (Version 2.3.0/1.53f)	ImageJ	Schindelin et al. ⁵⁷
MtrackJ	FIJI	Meijering et al. ⁵⁸
Python code for kinetochore oscillations analysis	This paper	https://github.com/miquelrosassalvans/kinetochore-oscillations-analysis.git
Other		
lipofectamine siRNAmix	Thermo Fisher Scientific	13778075
ProLong Gold Antifade reagent	Thermo Fisher Scientific	P36934

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the lead contact, Sophie Dumont (sophie.dumont@ucsf.edu).

Materials availability

This study did not generate new unique reagents.

Data and code availability

All data reported in this paper has been deposited at Mendeley Data and is publicly available as of the date of publication (<https://doi.org/10.17632/9gwpmj4hpr.1>).

All original code has been deposited at [github.com](https://github.com/miquelrosassalvans/kinetochore-oscillations-analysis.git) and is publicly available as of the date of publication (<https://github.com/miquelrosassalvans/kinetochore-oscillations-analysis.git>).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Rpe1-GFP cells (gift from A. Khodjakov, Wadsworth Center⁴⁰) were cultured in DMEM/F12 (Dulbecco's Modified Eagle Medium: Nutrient Mixture F-12, Thermo Fisher Scientific, 11320082) supplemented with 10% qualified and heat-inactivated fetal bovine serum (FBS) (10438-026, Gibco) and penicillin/streptomycin and maintained at 37°C and 5% CO₂. Cells used in this study were not authenticated by the authors.

METHOD DETAILS

Cell culture, siRNA transfection and drug treatments in Rpe1 cells

Cells were plated in 35 mm glass-bottom dishes (poly-D-lysine coated; MatTek Corporation) for live imaging experiments or in six wells plates after addition on #1.5 25 mm coverslips (acid cleaned and poly-L-lysine coated) for immunofluorescence experiments. For knockdown experiments, siRNA targeting SKAP (siSKAP: 5'-AGGCUACAAACCACUGAGUAA-3'³¹; siSKAP2: 5'-GAAAGA GUCCGAUUCUAG-3'³⁰) or luciferase siRNA (5' CGUACGCGGAUACUUCGA 3', control, WD01818022) were transfected in Rpe1-GFP cells and, using 4 μl of lipofectamine siRNAmix (Thermo Fisher Scientific, 13778075) and 10 μM of siRNA for 2 ml of cell culture media. Cells were incubated at 37°C for 6–8 h before media wash, and imaged or processed 24 h after transfection. SiR-tubulin (Cytoskeleton, CY-SC002) was used at 1 μM concentration and added to the cells 40 min before imaging in experiments for [Figure 2C](#). Nocodazole (Sigma-Aldrich, M1404-50MG) was used at 2 μM concentration and added 3 h before fixation in experiments for [Figures 2D](#) and [2E](#). MG132 (carbobenzoxy-Leu-Leu-leucinal, EMB Millipore, 474790-5MG) was used at 10 μM concentration in experiments for [Figure S2](#).

Immunofluorescence and immunoblotting

To validate the SKAP siRNA ([Figures S1A](#) and [S1B](#)), cells were seeded in six well plates, transfected with luciferase (control) or SKAP siRNA 24 h after transfection and processed for immunoblotting. Cells were collected in PBS1X (Phosphate Buffered Saline solution) using a cell scraper and lysed in PBS1X + 1% NP40 on ice for 30 min. Samples were run in a 4–12% Bis-Tris gel (Invitrogen, NPO335BOX) and transferred to a nitrocellulose membrane (Thermo Scientific, 88018). The following primary and secondary antibodies and dyes were used (incubated in TBS1X (Tris-buffered saline), 3% milk, 0.1% Tween for 1 h or 45 min, respectively): anti-SKAP (1 μg/ml, rabbit, Origene, TA333584), anti- α -tubulin (DM1A, 1:1000, mouse, Sigma, T6199), goat anti-mouse IgG-HRP (1:1000, sc-2005, Santa Cruz Biotechnology) and mouse anti-rabbit IgG-HRP (1:1000, sc-2357, Santa Cruz Biotechnology). Blots were exposed with SuperSignal West Pico Substrate (Thermo Scientific) and imaged with a Bio-Rad ChemiDoc XRS+ system.

For immunofluorescence experiments cells were fixed in 99.8% methanol for 10 min at -20°C, 24 h after siRNA transfection, and permeabilized in PBS1x, 0.5% BSA, 0.1% Triton (IF buffer thereafter) for 30 min. The following primary antibodies were incubated for 1 h in IF buffer: SKAP (1 μg/ml, rabbit, gift from I. Cheeseman⁵⁵), Kif18A (2 μg/ml, rabbit, Bethyl, A301-080A-M), MCAK (1:500, rabbit, cytoskeleton, AKIN05), α -Tubulin DM1A (1:1000, mouse, Sigma, T6199), p-Hec1-s69 (1:3000, rabbit, gift from J. DeLuca²⁵), CREST (1:100, human, Antibodies Incorporated, 15-234-0001), Hec1 (1:200, mouse, Abcam, ab3613), tubulin (1:2000, rat, Bio-rad, MCA77G). Three washes in IF buffer (10 min each) were done before incubation with the following secondary antibodies (1:1000 in IF buffer, 45 min): goat anti-mouse IgG Alexa Fluor 405, 488 and 568 (A31553, A11001 and A11004, Invitrogen), goat anti-rabbit IgG Alexa Fluor 405 and 568 (A31556 and A11011, Invitrogen), goat anti-rat IgG Alexa Fluor 488 (A11006, Invitrogen) and goat anti-human IgG Alexa Fluor 568 (A21090, Invitrogen). Samples were washed once in IF buffer and twice in PBS1X (10 min each) before mounting in ProLong Gold Antifade reagent (P36934, Thermo Fisher).

Microscopy and laser ablation

Samples were imaged using an inverted microscope (Eclipse Ti-E; Nikon) with a spinning disk confocal (CSU-X1; Yokogawa Electric Corporation), head dichroic Semrock Di01-T405/488/568/647 for multicolor imaging, equipped with 405 nm (100 mW), 488 nm (120 mW), 561 nm (150 mW), and 642 nm (100 mW) diode lasers, emission filters ET455/50M, ET525/50M, ET630/75M and ET690/50M for multicolor imaging, and an iXon3 camera (Andor Technology) operated by MetaMorph (7.7.8.0; Molecular Devices).⁴⁶ Cells were imaged through a 100X 1.45 Ph3 oil objective and 1.5X lens.

For live imaging and laser ablation experiments, cells were maintained in a stage-top incubation chamber (Tokai Hit) at 37°C and 5% CO₂. Metaphase oscillations were imaged every 3 s ([Figures 1](#), [2B](#), and [S2](#)). Laser ablation (30–40 pulses of 3 ns at 20 Hz) with 514 nm light was performed using the MicroPoint Laser System (Andor). For laser ablation experiments, images were acquired more slowly prior to ablation and then acquired more rapidly after ablation (3 s prior and 0.5 s after k-fiber ablation, and 6 s prior and 3 s after kinetochore ablation [[Figures 3](#) and [4](#), respectively]). Successful k-fiber ablation was verified by immediate K-K distance relaxation ([Figures 2G–2I](#), [3](#), and [S3](#)) and posterior front kinetochore movement poleward by dynein pulling. Successful kinetochore ablation was verified by immediate poleward movement of the remaining sister kinetochore ([Figure 4](#)). For long term imaging experiments ([Figure 2C](#)), Rpe1-GFP cells treated with 100 nM SiR-DNA were imaged every 4 min for 18–20 h using a 20X objective.

Study design and data inclusion criteria

Two general criteria for inclusion of cells in metaphase oscillation and laser ablation experiments were applied. First, cells must express detectable levels of GFP-CenpA at kinetochores, but not so high as to completely label chromosome arms. Second, cells must be in metaphase, with a defined metaphase plate. For oscillation experiments (Figures 1, 2B, and S2), 2-4 kinetochore pairs per cell were analyzed, and the two kinetochores from the pair must stay in focus for a minimum of 90 s. For MG132 treatment experiments (Figure S2), cells with low centromere stiffness were selected if their mean K-K distance over time was higher than two standard deviations over the mean control K-K distance. For kinetochore speed calculations in ablation experiments (Figures 3E and 4D), kinetochore tracks shorter than 5 timepoints were excluded due to the absence of a consistent directional movement.

We did not pre-estimate a required sample size before performing experiments nor did we blind or randomize samples during experimentation or analysis. The ablation experiments in this study are low throughput by nature, which does not enable us to report averages from multiple independent replicate experiments. Instead, we pool cells from across different independent experiments (with at least three independent experiments per condition per assay).

QUANTIFICATION AND STATISTICAL ANALYSIS

Analyzing fluorescence intensity

Images from immunofluorescence experiments were processed and fluorescence intensity was quantified using FIJI (Version 2.3.0/1.53f).⁵⁷ For protein intensity measurements, a color threshold mask (Yen method) was applied using the CREST or Hec1 signal (for kinetochore selection) or DM1A signal (for spindle microtubule selection) to define the areas in which the fluorescence intensity would be measured for each protein of interest. Fluorescence intensity was measured for the proteins of interest (Hec1-S69, MCAK, Kif18A) and for the reference (CREST, Hec1 or Tubulin). Intensity of the protein of interest was normalized by the kinetochore or microtubule marker intensity by dividing the total intensity of the protein of interest by the intensity of the reference in the selected area. Figures 3H, S4C, and S4D showing data from individual representative experiments, three independent experiments were performed for each quantification, obtaining comparable results.

Analyzing kinetochore behaviors

Kinetochores and centrioles were manually tracked from GFP-CenpA /centrin1-GFP videos using the MtrackJ plugin from FIJI.⁵⁸ GFP-centriole position was used as a marker for the spindle pole position. Kinetochore position was calculated as the distance from the spindle pole position. In metaphase oscillations experiments, all quantifications and statistical analyses were performed using home-written Python code (<https://github.com/miquelrosassalvans/kinetochore-oscillations-analysis.git>). Kinetochore speed at each timepoint was calculated as the difference in kinetochore position between two consecutive timepoints. Kinetochore speed was calculated by obtaining the slope of the best fitting regression line of individual kinetochore tracks (Figures 3E and 4D) or of entire tracks together (Figures 3C, 3D, and 4C). Sister kinetochore movement coordination was obtained by calculating the correlation of sister kinetochore velocity over time or by the percentage of timepoints in which sister kinetochore movement direction was opposite. Kinetochore directional switch was determined by the consistent movement of a kinetochore in a the new direction for 3 consecutive timepoints. In ablation experiments, time after ablation was measured from the first timepoint immediately after ablation. In all experiments, a kinetochore directional switch was defined by the consistent movement of the kinetochore in the opposite direction for a period of 3 consecutive timepoints (9 s in oscillations and kinetochore ablations and 1.5 s for k-fiber ablation experiments). K-K distance was calculated by subtracting the position of sister kinetochore A from sister kinetochore B, obtaining the length of the vector, and calculating the K-K distance average over time for each kinetochore pair. K-K distance relaxation was obtained by subtracting the K-K distance at the first timepoint after ablation from the last timepoint before ablation.

Statistical analysis

Statistical analysis was performed in Python or Graphpad (Prism 9). The Fisher's exact test was used in Figure 1G, Student's t test was used for parametric datasets (Figures 2B, 2I, 3E, 3H, 4E, S4C, and S4D), Mann-Whitney test for non-parametric datasets (Figures 1C-1F, 2C, 2E, 3F, 4D, S2A, S2C, and S2D) and analysis of covariance test, ANCOVA, for linear regression slopes comparison (Figures 3C, 3D, 4C, and S3).

Video preparation

Videos were prepared using FIJI. Brightness and contrast were linearly adjusted to clearly visualize the kinetochores and centrioles.